

## **4376 FIRST AID KIT- 4376 first aid kit**

**Honeywell Safety Products USA, INC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

## **4376 First Aid Kit (BZK wipe,FABC, neomycin, alcohol wipes, sting relief, ASA, aypanal-Z019807)**

### **Alcohol Wipe**

#### ***Active ingredient***

Isopropyl alcohol 70%

### **Alcohol Wipe**

#### ***Purpose***

First aid antiseptic

### **Alcohol Wipe**

#### ***Uses***

- first aid to help prevent infection in minor cuts, scrapes, and burns

### **Alcohol Wipe**

#### ***Warnings***

#### **For external use only**

#### **Do not use**

- in the eyes
- over large areas of the body

#### **Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burn

#### **When using this product**

- do not use longer than one week unless directed by a doctor

#### **Stop use and consult a doctor**

- if condition persists or gets worse

#### **Keep out of reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away.

### **Alcohol Wipe**

#### ***Directions***

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

## **Alcohol Wipe**

### ***Other information***

store at room temperature 15 ° to 25 ° C (59 ° to 77 °F)

## **Alcohol Wipe**

### ***Inactive ingredient***

water

## **Alcohol Wipe**

### ***Questions***

1-800-430-5490

## **Aspirin**

### ***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\* \*nonsteroidal anti-inflammatory drug

## **Aspirin**

### ***Purpose***

Pain reliever/fever reducer

## **Aspirin**

### ***Uses***

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache
- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

## **Aspirin**

### ***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling

- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:are:

- age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout or arthritis

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- ringing in the ears or loss of hearing occurs
- any new symptoms appear

**If pregnant or breast-feeding,**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

- In case of overdose, get medical help or contact Poison Control Center right away.

**Aspirin**

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last,

not more than 12 tablets in 24 hours

- children under 12 years of age: consult a doctor

## **Aspirin**

### ***Other information***

- store at room temperature 15° - 30°C (59° - 86°F)
- TAMPER EVIDENT PACKETS
- DO NOT USE IF OPEN OR TORN

## **Aspirin**

### ***Inactive ingredients***

corn starch, croscarmellose sodium\*, hypromellose\*, microcrystalline cellulose\*, mineral oil\*, polyethylene glycol\*, povidone, propylene glycol, silicon dioxide, stearic acid\*, titanium dioxide\*

\*may contain these ingredients

## **Aspirin**

### ***Questions or Comments***

1-800-430-5490

## **Sting Relief**

### ***Active ingredient***

Ethyl alcohol 50.0%

Lidocaine HCl 2.0%

## **Sting Relief**

### ***Purpose***

Antiseptic

Topical pain relief

## **Sting Relief**

### ***Uses***

- prevent infection in minor scrapes, and temporary relief of itching of insect bites

## **Sting Relief**

### ***Warnings***

**For external use only**

**Flammable, keep away from open fire or flame**

### **Do not use**

- over large areas of the body
- in eyes
- over raw or blistered areas

### **Stop use and ask a doctor**

- if conditions worsen or persist for more than 7 days or clear up and occur again within a few days

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Sting Relief*****Directions***

- adults and children 2 years and older: Apply to cleaned affected area not more than 3 times daily.
- children under 2 years of age: consult a doctor.

**Sting Relief*****Inactive ingredients***

benzalkonium chloride, menthol, and purified water

**Sting Relief*****Questions or Comments?***

1-800-430-5490

**BZK*****Active ingredient***

Benzalkonium chloride 0.13% w/v

**BZK*****Purpose***

First aid antiseptic

**BZK*****Uses***

Antiseptic cleansing of face, hands, and body without soap and water

**BZK*****Warnings*****For external use only****Do not use**

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

**Stop use and ask a doctor if**

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **BZK**

### ***Directions***

- tear open packet and use as a washcloth

## **BZK**

### ***Other information***

- store at room temperature 15 ° to 30 ° C (59 ° - 86 ° F)
- do not reuse towelette

## **BZK**

### ***Inactive ingredient***

water

## **BZK**

### ***Questions***

1-800-430-5490

## **Neomycin**

### ***Active ingredient (each gram contains)***

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

## **Neomycin**

### ***Purpose***

First aid antibiotic

## **Neomycin**

### ***Uses***

- first aid to help prevent infection in
- minor cuts
- scrapes
- burns

### ***Do not use***

- in the eyes
- over large areas of the body

## **Neomycin**

### ***Warnings***

### ***For external use only***

### ***Ask a doctor before use if you have***

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away

**Neomycin*****Directions***

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

**Neomycin*****Other information***

- store at 15 ° to 25 ° C (59 ° to 77 ° F)

**Neomycin*****Inactive ingredient***

petrolatum

**Neomycin*****Questions***

1-800-430-5490

**Aypanal EX*****Active ingredient***

Acetaminophen 500 mg

**Aypanal EX*****Purpose***

Pain reliever/fever reducer

**Aypanal EX*****Uses***

- temporarily relieves minor aches and pains due to the common cold and headache - temporarily reduces fever

**Aypanal EX*****Warnings***

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount.

- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away

### **Do Not Use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if you have**

- liver disease

### **Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- If pregnant or breastfeeding,
- ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Aypanal EX**

### **Directions**

- **do not take more than directed (see overdose warning)**
- adults and children 12 years of age and over: Take 2 tablets with water every 6 hours while symptoms last.
- do not take any more than 8 tablets in 24 hours.
- children under 12: consult a doctor

## **Aypanal EX**

### **Other information**

- store at room temperature 15 ° -30 ° C (59 ° -86 ° F)
- TAMPER EVIDENT- DO NOT USE IF OPEN OR TORN

## **Aypanal EX**

### **Inactive ingredients**

microcrystalline cellulose, povidone, sodium starch glycolate, starch, stearic acid



## **Aypanal EX**

### ***Questions or Comments?***

1-800-430-5490

## **FABC**

### ***Active ingredient***

Benzalkonium chloride 0.13%

Lidocaine HCl 0.5%

## **FABC**

### ***Purpose***

First Aid antiseptic

External analgesic

## **FABC**

### ***Uses***

- prevent skin infection
- for temporary relief of pain associated with minor burns

## **FABC**

### ***Warnings***

For external use only

### **Do not use**

- in or near the eyes
- if you are allergic to any of the ingredients
- in large areas of the body, particularly over raw surfaces or blistered areas
- for more than 10 days

### **Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

### **Stop use and ask a doctor if**

- condition worsens
- symptoms persist for more than 7 days or clear up and occurs again within a few days

### **Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **FABC**

### ***Directions***

- **adults and children 2 years of age and older:**
- clean the affected area
- apply a small amount of this product (equal to the surface area of the tip of a finger) onto affected

area 1 to 3 times daily

- may be covered with a sterile bandage
- **children under 2 years of age: consult a doctor**

## **FABC**

### ***Other information***

- tamper evident sealed packets
- do not use if packet is opened or torn

## **FABC**

### ***Inactive ingredients***

aloe barbadensis juice, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl stearate SE, methylparaben, mineral oil, PEG-100, propylene glycol, propylparaben, stearic acid, trolamine, water

## **FABC**

### ***Questions***

1-800-430-5490

## **4376**


### **Z019807 KIT CONTENTS**

1 ADHESIVE TAPE W/P 1/2"X 5 YD  
1 FIRST AID GUIDE ASHI  
1 EMERGENCY SURVIVAL BLANKET  
1 GAUZE CLEAN-WRAP BDGE N/S 2"  
10 CTA 6" SINGLE TIP 100/BAG  
1 SCISSOR BDGE 4" RED PLS HDL  
1 FANNY PACK RED FAK LOGO EMPTY  
4 TONGUE BLADES SR WRAPPED 250  
LBL STOCK 6-3/8"X4"  
1 LBL STOCK 3"x1-7/8"  
1 LBL CONTS 6 3/4"X3 1/2" ID B  
6 BZK ANTISEPTIC WIPE, BULK  
1 PR LRG NITRILE GLVES ZIP BAG  
3 FIRST AID BURN CREAM 1.0GR PKT EACH  
3 POUCH NEOMYCIN ANTIBIOTIC .9 G  
4 ADH BANDAGE BUTTERFLY 1980000  
6 WIPE ALCOHOL PREP IPA 70% (DUKAL)  
2 LANCET BLADE #11 STERILE  
10 ADH BANDAGE 3/8" X 1 1/2" DNX  
3 SAFETEC STING RELIEF WIPES BULK

1 COLD PACK UNIT 4"X6" BULK  
 4 GAUZE PADS 2"X2" 12PLY  
 2 GAUZE PADS 4"X4" 12PLY  
 15 PLASTIC BANDAGE 3/4" X 3"  
 1 WOVEN KNUCKLE BANDAGE  
 1 ZIP-LOCK BAG 5" X 5" .002  
 2 AYPANAL BULK 2/PK  
 2 ASPIRIN BULK 2/PK

## Alcohol Wipe Label

825362T Rev. A  
 white printed  
 Pms 072 blue, Pms 186 red,  
 Pms 7459 blue and black



## Aspirin Principal Display Panel



47001083  
Rev B

Antiseptic Towelettes

**Honeywell**

02-16-35MD

## Antiseptic Towelettes

*Benzalkonium chloride*  
*First aid antiseptic*

**Six-Saturated Towelettes**

Distributed by  
Honeywell Safety  
Products USA, Inc.  
Smithfield, RI 02917

## Antiseptic Towelettes

Honeywell

47001083  
Rev B

### Drug Facts

#### Active Ingredient

Benzalkonium chloride 0.133% w/v

#### Purpose

First aid antiseptic

#### Uses

- antiseptic cleaning of face, hands and body without soap and water.
- air dries in seconds

#### Warnings

##### For external use only

**When using this product** • do not use in the eyes or apply over large areas of the body

**Ask a doctor before use** • in case of deep or puncture wounds, animal bites, or serious burns

#### Stop use and consult a doctor if

- irritation, redness or other symptoms develop
- condition persists or gets worse

**Do not use** • longer than 1 week unless directed by doctor

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

- tear open packet, unfold and use as washcloth

#### Other Information



- store at room temperature 15° -30° C (59° -86° F)
- do not reuse towelette

**Inactive ingredient** water


**Questions or comments** 1-800-430-5490

**Neomycin**  
**Principal Display Panel**

796041-25 Rev A Unit Carton Printing Plate for "C" size carton.



Principal Display Panel



FABC  
Principal Display Panel



901850-10	First Aid Burn Cream	021020-10
		<p><b>First Aid Burn Cream</b> Benzalkonium chloride First aid antiseptic Lidocaine HCl External analgesic</p> <p>Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917</p> <p>10 Packets, Net Wt 1/32 oz (0.9g) each</p>

901850-10	First Aid Burn Cream	<p><b>Drug Facts</b></p> <table border="1"> <tr> <th>Active ingredients</th> <th>Purposes</th> </tr> <tr> <td>Benzalkonium chloride 0.13%</td> <td>First aid antiseptic</td> </tr> <tr> <td>Lidocaine HCl 0.5%</td> <td>External analgesic</td> </tr> </table> <p><b>Use</b> - prevent skin infection - for the temporary relief of pain associated with minor burns</p> <p><b>Warnings For external use only</b></p> <p><b>Do not use</b> - in or near the eyes - if you are allergic to any of the ingredients - in large areas of the body particularly over raw surfaces or blistered areas - for more than 10 days</p> <p><b>Ask a doctor before use if you have</b> - deep or puncture wounds - animal bites - serious burns</p> <p><b>Stop use and ask a doctor if</b> - condition worsens - symptoms persist more than 7 days or clear up and occur again within a few days</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away</p> <p><b>Directions</b> <b>adults and children 2 years and over:</b> - clean the affected area - apply a small amount of this product (equal to the surface area of the tip of a finger) onto affected area 1 to 2 times daily - may be covered with a sterile bandage <b>children under 2 years:</b> ask a doctor</p> <p><b>Other information</b> - tamper evident sealed packets - do not use if packet is opened or torn</p> <p><b>Inactive ingredients</b> aloe barbadensis leaf juice, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl stearate SE, methylparaben, mineral oil, PEG-100, propylene glycol, propylparaben, stearic acid, triethanolamine, water</p> <p><b>Questions or comments?</b> 1-800-430-5430</p>	Active ingredients	Purposes	Benzalkonium chloride 0.13%	First aid antiseptic	Lidocaine HCl 0.5%	External analgesic
		Active ingredients	Purposes					
Benzalkonium chloride 0.13%	First aid antiseptic							
Lidocaine HCl 0.5%	External analgesic							

4376 Kit Label  
Z019807



# FIRST AID KIT



Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

APPROVED

By Rodrigo Rosas Asilano at 2:58 pm, Mar 11, 2019

## 4376 FIRST AID KIT

4376 first aid kit kit

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4376
--------------	----------------	--------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4376-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	3 PACKET	2.7 g
Part 2	6 POUCH	2.4 mL
Part 3	2 PACKET	4
Part 4	3 PACKET	2.7 g
Part 5	6 PACKET	8.4 mL
Part 6	3 POUCH	1.2 mL
Part 7	2 PACKET	4

Part 1 of 7

FIRST AID BURN

benzalkonium chloride, lidocaine hydrochloride cream

Product Information

Item Code (Source) NDC:0498-0903

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE	0.5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
PEG-100 STEARATE (UNII: YD01N1999R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
TROLAMINE (UNII: 9O3K93S3TK)	
WATER (UNII: 059QF0KO0R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	12/20/2017	

Part 2 of 7

ALCOHOL WIPE

isopropyl alcohol swab

Product Information

Item Code (Source)	NDC:0498-0143
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	09/18/2018	

Part 3 of 7

ASPIRIN

aspirin tablet

Product Information

Item Code (Source)	NDC:0498-0114
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/18/2018	

Part 4 of 7

NEOMYCIN

antibiotic ointment

Product Information

Item Code (Source)	NDC:0498-0730
Route of Administration	TOPICAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g

Inactive Ingredients	
Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0730-01	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	03/31/2010	

Part 5 of 7	
ANTISEPTIC TOWELETTE	
benzalkonium chloride liquid	

Product Information	
Item Code (Source)	NDC:0498-0501
Route of Administration	TOPICAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0501-00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	12/22/2017	

Part 6 of 7

STING RELIEF PAD

ethyl alcohol, lidocaine swab

Product Information

Item Code (Source)	NDC:0498-0733
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	20 mg in 1 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	0.5 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
MENTHOL (UNII: L7T10EP3A)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0733-00	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	12/23/2017	

Part 7 of 7

AYPANAL EX

acetaminophen tablet

## Product Information

**Item Code (Source)** NDC:0498-2110

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	FR1
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-2110-01	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/02/2017	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

**Labeler** - Honeywell Safety Products USA, INC (079287321)

Establishment			
Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, INC		079287321	pack(0498-4376)

Establishment			
Name	Address	ID/FEI	Business Operations
Ultra Seal Corporation		085752004	manufacture(0498-0114, 0498-2110)

Establishment			
Name	Address	ID/FEI	Business Operations
Water-Jel Technologies		155522589	manufacture(0498-0730, 0498-0903)

Establishment			
Name	Address	ID/FEI	Business Operations
Changzhou Maokang Medical		421317073	manufacture(0498-0143, 0498-0501)

Establishment			
Name	Address	ID/FEI	Business Operations
Safetec of America Inc		874965262	manufacture(0498-0733)

Revised: 6/2019

Honeywell Safety Products USA, INC